

Unintentional Injury for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights

- Unintentional injury is the leading cause of death among youth.
- Unintentional injuries are caused by behaviors such as not wearing a seatbelt, riding a bike or motorcycle without a helmet, drinking and driving, riding with a drunk driver and not wearing a life jacket.
- Motor vehicle crashes are the most common type of injury death among youth.









Prevalence **	2002	2004	2006	2008	2010
Rode with drinking driver	S	25% ± 5% (342)	S	26% ± 4% (407)	20% ± 2 (1078)
Drove after drinking	S	11% ± 3% (340)	S	13% ± 3% (407)	9% ± 2° (1077)

From 2002 to 2004, change in riding with a drinking driver could not be assessed.



Most unintentional injuries are not accidents because they can be prevented. Alcohol and other substances impair one's judgment and may contribute to the likelihood of many injuries and even death. For more information and resources, visit the Washington State Injury and Violence Prevention Website at: http://www.doh.wa.gov/hsqa/emstrauma/injury/

^{*} indicates a significant change (Wald X², p<.05)

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location


Violent Behaviors for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- The federal Gun Free Schools Act requires the expulsion of students who bring firearms to school (SED. 4141) and Washington Law requires a one year expulsion for possession of firearms or dangerous weapons on school campuses. (RCW 28A.600.420 / RCW 9.41.280)
- Schools in Washington are required to have a comprehensive school safety plan that includes prevention, intervention, all-hazards/crisis response and post crisis recovery. (RCW 28A.320.125)








Violence and Academic Failure Trends

	Prevalence **	2002	2004	2006	2008	201
	In a physical fight at school	S	9% ± 3% (341)	S	7% ± 2% (407)	5% ± (532
	Academic failure	S	49% ± 5% (350)	S	45% ± 5% (408)	46% ± (549
	Carried a weapon at school	S	6% ± 2% (689)	S	8% ± 2% (815)	4% ± (1087



Schools must have in place effective referral systems that encourage students to report threats if they hope to prevent violent incidents.

For More Information

Supportive learning environments are essential for student academic achievement. Students need safe, nurturing, healthy, and civil schools in order to be successful.

For More Information about school safety, visit the OSPI-sponsored Safety Center Website at: http://www.K12.wa.us/safetycenter/

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net

Harassment, Intimidation, and Bullying for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- Washington State Law (RCW 28A.300.285) requires all school districts to adopt a policy that prohibits harassment, intimidation, and bullying (HIB).
- The legislation also requires the Office of Superintendent of Public Instruction (OSPI) to publish a model policy at the following website: http://www.k12.wa.us/Safetycenter/LawEnforcement/Student Discipline.aspx



Bullying is defined as a student, or group of students, saying or doing nasty or unpleasant things to another student. It is also bullying when a student is teased repeatedly in a way he or she doesn't like.

Bullied in Past Month

Bullying and Academic Failure Trends Grade 12 Bullied in past month Academic failure Feel safe at school 100% 80% 60% 40% 2002 2004 2006

Prevalence **	2002	2004	2006	2008	20
Bullied in past month	S	15% ± 3% (664)	S	16% ± 3% (770)	16% (10
Academic failure	S	49% ± 5% (350)	S	45% ± 5% (408)	46% (5
Feel safe at school	S	83% ± 3% (696)	S	84% ± 2% (819)	90% (108

Compared to the State, All Grades, 2004 Local State 100% 80% 60%

40% 20% 0% 8th 10th 12th 28% 33% 20% 15% 30% 29% 22% 16% Diff. 3% * -1% -1% -1%

Researchers have identified best practice support programs that, when implemented with fidelity in schools and classrooms, can address school harassment & bullying and build positive school cultures.



Supportive learning environments are essential for student academic achievement. Students need safe, nurturing, healthy, and civil schools in order to be successful.

For More Information visit the OSPI-sponsored Safety Center Website at: http://www.K12.wa.us/safetycenter/

 $^{^{\}star}$ indicates a significant change (Wald X^2 , p<.05)

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Community Risk Factors for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Noteworthy

- Research shows that the risk factors measured by the Healthy Youth Survey are linked to adolescent substance use, delinquency, violence, school dropout, and other risky health behaviors.
- The more risk factors a youth is exposed to, the more likely it is that she or he will engage in problem behaviors.

2004 Community Risk Factor Scales, % at Risk Grade 12







Community Risk Factor Trends, % at Risk
Grade 12

Pro drug-use laws and norms
Availability of drugs
Availability of handguns
Low neighborhood attachment

60%
50%
40%
30%
20%
10%
2002
2004
2004
2006

Prevalence **	2002	2004	2006	2008	201
Pro drug-use laws and norms	S	33% ± 5% (353)	S	32% ± 5% (410)	30% ±
Availability of drugs	S	45% ± 5% (354)	S	45% ± 5% (407)	33% ± (555
Availability of handguns	S	22% ± 4% (353)	S	18% ± 4% (406)	17% ±
Low neighborhood attachment	S	N/S	S	51% ± 5% (414)	48% ±

From 2002 to 2004, change in the perceived availability of drugs could not be assessed.

For More Information

Parents, schools, kids, and communities can work together to keep youth safe. To see how State and local partners are addressing these issues, see listings on the AskHYS homepage. For free guides on preventing alcohol and other drug use, call Washington State Alcohol/Drug Information Clearinghouse: 1-800-662-9111.

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location

Community Protective Factors for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:


Noteworthy

- Kids living in high-risk environments can be protected from problem behaviors by positive relationships with adults who care about their healthy development.
- Providing opportunities for involvement in positive and healthy activities help communities promote youth development.


of me

2004 Rewards for Prosocial Community Involvement Scale and Selected Items, % Protected Grade 12 100% 80% 52% 60% 40% 26% 20% 0% Protective Neighbors Neighbors Neighbors Factor encourage proud Scale my good my best

2004 Opportunities for Prosocial Community Involvement Scale and Selected Items Grade 12




Rewards for Prosocial Community Involvement Compared to the State, All Grades, 2004



Protective Factor Trends, % Protected Grade 12

Rewards for Prosocial Community Involvement
Opportunities for Prosocial Community Involvement



Prevalence **	2002	2004	2006	2008	20
Rewards for social Community Involvement	S	52% ± 5% (353)	S	60% ± 5% (412)	٨
Opportunities for isocial Community Involvement	S	69% ± 5% (348)	S	77% ± 4% (408)	77% (5

From 2002 to 2004, change in community opportunities for prosocial involvement could not be assessed.



For More Information

Parents, schools, kids, and communities can work together to keep youth safe. To see how State and local partners are addressing these issues, see listings on the AskHYS homepage. For free guides on preventing alcohol and other drug use, call Washington State Alcohol/Drug Information Clearinghouse: 1-800-662-9111.

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


School Risk Factors for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights

- Research has identified a unique set of risk factors that put youth at risk for problems, such as substance use and school drop out.
- Youth who get poor grades, have friends who smoke, experience family conflict, and live in neighborhoods that are disorganized are at greater risk for negative youth behaviors.









Prevalence **	2002	2004	2006	2008	201
Current alcohol use	S	41% ± 4% (681)	S	45% ± 3% (795)	38% : (108
Academic failure	S	49% ± 5% (350)	S	45% ± 5% (408)	46% : (54
Low commitment to school	S	43% ± 5% (356)	S	41% ± 5% (413)	36% : (55

Students experiencing academic failure and lack of commitment to school are at higher risk for substance abuse, delinquency, violence, pregnancy and school drop out.



Supportive learning environments are essential for student academic achievement. Students need safe, nurturing, healthy, and civil schools in order to be successful. Researchers have identified best practice support programs that, when implemented with fidelity in schools and classrooms, can decrease risk factors and build positive school cultures.

For More Information about school risk factors, visit the website http://captus.samhsa.gov/western

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net

School Protective Factors for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights


- Research has identified a unique set of protective factors that buffer youth from problems such as substance use and school drop out.
- Youth who are involved in positive school activities, have well developed social skills, are attached to their families, and live in neighborhoods that invite positive participation are at less risk for negative youth behaviors.

2004 Elements of School Protection Grade 12 100% 86% 83% 80% 50% 60% 40% 27% 20% 0% Teachers **Parents** Opportunity Chances notified to talk to involved


hard of good teacher involved work

Offering young people opportunities for meaningful involvement in school and recognizing them for their efforts and accomplishments will help buffer them from problem behavior.

Element of Protection - Feeling Safe at School Compared to the State, All Grades, 2004



Protection Factors and Alcohol Use Trends Grade 12



Prevalence **	2002	2004	2006	2008	20:
Current alcohol use	S	41% ± 4% (681)	S	45% ± 3% (795)	38% : (108
Opportunities for school involvement	S	63% ± 5% (352)	S	65% ± 5% (411)	67% : (55
Rewards for school involvement	S	44% ± 5% (351)	S	51% ± 5% (412)	43% : (55

Students in schools with low levels of protection and even moderate substance use are less likely to meet the standards for the mathematics, reading and writing sections of standardized tests.



Supportive learning environments are essential for student academic achievement. Students need safe, nurturing, healthy, and civil schools in order to be successful. Researchers have identified best practice support programs that, when implemented with fidelity in schools and classrooms, can increase protective factors and build supportive school cultures.

For More Information about school protective factors, visit the website http://captus.samhsa.gov/western

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Peer-Individual Risk Factors for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights

- Research shows that the risk factors measured by the Healthy Youth Survey are linked to adolescent substance use, delinquency, violence, school dropout, and other risky health behaviors.
- The attitudes kids have about drug use strongly influence the likelihood they will use drugs. Families and communities should model healthy attitudes and choices.


2004 Peer-Individual Risk Factor Scales, % at Risk Grade 12 60% 44% 50% 34% 40% 30% 30% 20% 10% 0% Perceived Attitudes Friends Intentions risk of favor use to use drugs drug use drug use druas







Prevalence **	2002	2004	2006	2008	2010
Perceived risk of drug use	S	44% ± 5% (339)	S	46% ± 5% (394)	48% ± 4 (551)
Attitudes favor drug use	S	34% ± 5% (317)	S	41% ± 5% (372)	36% ± 4 (546)
Friends use drugs	S	31% ± 5% (306)	S	34% ± 5% (360)	24% ± 4 (540)
Intentions to use drugs	S	30% ± 5% (305)	S	38% ± 5% (358)	34% ± 4 (540)



For More Information

Parents, schools, kids, and communities can work together to keep youth safe. To see how State and local partners are addressing these issues, see listings on the AskHYS homepage. For free guides on preventing alcohol and other drug use, call Washington State Alcohol/Drug Information Clearinghouse: 1-800-662-9111.

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location

Peer-Individual Protective Factors for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights


• Kids are strongly influenced by their peers. Parents need to know who their children's friends are, where they are and what they are doing.


0%

2002

Kids who have the skills and opportunities to be involved in pro-social activities are less likely to engage in risky behavior.

2004 Peer-Individual Protective Factor Scales, % Protected Grade 12 100% 80% 68% 55% 60% 46% 42% 40% 20% 0% Prosocial Belief Prosocial Social in a involvement skills interaction moral






Peer-Individual Protective Factor Trends, % Protected Grade 12 Prosocial peer interaction Belief in a moral order Prosocial involvement Social skills 100% 80% 40% 20%

			,		
Prevalence **	2002	2004	2006	2008	2010
Prosocial peer interaction	N/S	46% ± 6% (310)	S	47% ± 5% (365)	54% ± 4 (540)
Belief in a moral order	S	55% ± 5% (344)	S	49% ± 5% (401)	55% ± 4 (554)
Prosocial involvement	N/S	42% ± 5% (354)	S	N/A	N/A
Social skills	S	68% ± 5% (341)	S	63% ± 5% (393)	50% ± 4 (530)

2004

2006



For More Information

Parents, schools, kids, and communities can work together to keep youth safe. To see how State and local partners are addressing these issues, see listings on the AskHYS homepage. For free guides on preventing alcohol and other drug use, call Washington State Alcohol/Drug Information Clearinghouse: 1-800-662-9111.

^{*} indicates a significant change (Wald X², p<.05)

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Family Protective Factors for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- Children who are attached to positive families are less likely to develop problems in adolescence.
- When parents set clear standards for children's behavior, when they are widely and consistently supported, and when the consequences for not following the standards are consistent, young people are more likely to follow the standards.


2004 Opportunities for Prosocial Family Involvement, % Protected Scale and Selected Items



In 2004, .% of 12th grade students in our school district were protected with opportunities for


prosocial family involvement.

Rewards for Prosocial Family Involvement Compared to the State, All Grades, 2004



Family Protective Factor Trends Grade 12

Rewards for prosocial family involvement Opportunities for Prosocial Family Involvement



Prevalence **	2002	2004	2006	2008	20
Rewards for prosocial family involvement	S	N/A	S	54% ± 5% (350)	46% (42
Opportunities for Prosocial Family Involvement	S	N/A	S	51% ± 5% (353)	50% (4

From 2002 to 2004, change in family opportunities for prosocial involvement could not be assessed.



For More Information

Parents and caregivers are the #1 influence on a child's decisions about alcohol and other drugs. You have the power to help your child make healthy choices.

Talking with your child about his or her activities opens up an opportunity for you to share your interests and values. Young people are much less likely to have mental health and substance abuse problems when they have positive activities to do and when caring adults are involved in their lives.

For More Information, visit the Start Talking Now website at www.StartTalkingNow.org.

^{*} indicates a significant change (Wald X², p<.05)


^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses. Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location


Weight and Obesity for Edmonds School District

Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights


- Youth who have fewer opportunities for physical activity and increased time spent in sedentary activities such as watching television and using computers are more likely to be overweight.
- Factors that contribute to overweight among youth include eating fewer meals at home, viewing more food advertisements targeting youth, and increased food portion sizes.






Prevalence **	2002	2004	2006	2008	2010
Overweight	S	8% ± 3% (318)	S	11% ± 3% (377)	10% ± (493
At risk for being overweight	S	14% ± 4% (318)	S	13% ± 3% (377)	13% ± (493
Not overweight	S	77% ± 5% (318)	S	76% ± 4% (377)	78% ± (493







For More Information

To be successful in reducing overweight among youth, governments, communities and people need to work together to create environments that support healthy diets and opportunities for physical activity.

For More Information and resources visit the Washington State Nutrition and Physical Activity Website at: www.doh.wa.gov/cfh/nutritionpa

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Dietary Behaviors for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- A nutritious diet can reduce major risk factors for chronic disease such as obesity, high blood pressure, and high blood cholesterol.
- Poor nutrition has the potential to affect the growth, development, and health status of children and adolescents.







	Youth Die	etary Behavior Trends Grade 12	
	* * * * * * * * * * * * * * * * * * *	Eat 5+ fruits/vegs. daily Usually eat family dinners Drink 2+ sodas daily	
100% -		<u> </u>	
80% -			
60% -			
40% -		A	
20% -		•	
0% -	2002	2004	2006

Prevalence **	2002	2004	2006	2008	2010
Eat 5+ fruits/ vegs. daily	S	20% ± 4% (323)	S	20% ± 4% (363)	N/S
Usually eat family dinners	S	46% ± 5% (328)	S	43% ± 5% (400)	51% ± 4 (527) *
Drink 2+ sodas daily	S	22% ± 5% (326)	S	13% ± 3% (397)	10% ± 3 (526)

From 2002 to 2004, change in usually eating dinner with families could not be assessed.

For More Information

Individual food choices are made within the context of the family and the community. In order to be able to choose a nutritious diet, an affordable supply of health-promoting foods must be available.

For More Information and resources visit the Washington State Nutrition and Physical Activity Website at: www.doh.wa.gov/cfh/nutritionpa

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Physical Activity for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights

- Regular physical activity reduces rates of obesity and serious diseases, helps people maintain a healthy body weight and improves quality of life
- The Centers for Disease Control and Prevention (CDC) recommends 60 minutes of physical activity on five or more days a week.
- One challenge to being more physically active is the amount of recreational screen time spent on sedentary activities such as watching TV, playing video games or using a computer for fun.









Prevalence **	2002	2004	2006	2008	201
Met mod/vigorous activity recs	S	66% ± 5% (322)	S	66% ± 5% (396)	N/S
2 or less hrs screen time daily	S	47% ± 5% (320)	S	45% ± 5% (392)	39% ± (523
Participate in PE classes daily	S	22% ± 5% (321)	S	16% ± 4% (396)	18% ±

From 2002 to 2004, change in 2 hours or less of daily screen time could not be assessed.

For More Information

Many social and environmental factors influence the decision to be active. Successful strategies to promote physical activity include educating individuals and developing policies to create environments that support being physically active.

For More Information and resources visit the Washington State Nutrition and Physical Activity Website at: www.doh.wa.gov/cfh/nutritionpa

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Asthma for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- In Washington State, about one in five youth have been diagnosed with asthma.
- Among youth with asthma, about one in five had at least one visit to an emergency department for asthma attack during 2006.
- Youth who smoke are more likely to have asthma.







Youth Asthma Trends Grade 12 Doctor diagnosed asthma Asthma attack or meds past year 40% 30% 10% 2002 2004 2006

Prevalence **	2002	2004	2006	2008	201
Doctor diagnosed asthma	S	17% ± 4% (318)	S	20% ± 4% (391)	21% ± (528
Asthma attack or meds past year	S	25% ± 5% (319)	S	N/S	N/S

All youth with asthma should have a written asthma action plan to follow in the event of an attack.



Asthma is a lung disease that causes trouble with breathing, usually in episodes called attacks. An attack can be triggered by exercise, exposure to smoke, mold, allergens like pet dander, and air pollution such as diesel exhaust. Asthma can be controlled by avoidance of asthma triggers, regular health care visits, compliance with medication schedules, and maintaining appropriate weight.

For More Information visit the Asthma Program's website at: http://www.doh.wa.gov/cfh/asthma

Washington Asthma Initiative's website at: http://www.alaw.org/asthma/washington_asthma_initiative/

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Depression & Suicide for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:


Highlights

- Suicide is the second leading cause of death for Washington teens 15-19 years old.
- Most youth suicides occur at home.










Prevalence **	2002	2004	2006	2008	201
Depression	S	36% ± 4% (645)	S	31% ± 3% (763)	29% ± (107
Considered attempting suicide	S	16% ± 4% (340)	S	14% ± 2% (762)	16% ± (107
Likely to seek help if depressed	S	58% ± 7% (206)	S	65% ± 6% (262)	59% ±



For More Information

Depression is a feeling of sadness, helplessness and hopelessness that lingers over time. Untreated it can lead to alcohol abuse and even suicide. For more information and resources, visit the Washington State Youth Suicide Prevention Program Website at:

http://www.yspp.org

National Suicide Prevention Hotline

- phone number 1-800-273 TALK(8255)
- website: http://www.suicidepreventionlifeline.org

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.


Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location Report generated at www.AskHYS.net


Current Substance Use for Edmonds School District


Year: 2004 Grade: 12 Gender: Both Ethnicity:

Highlights

- Parents are the primary influence on their children's decision to use alcohol, tobacco, and other drugs.
- Brain development continues into the early twenties. The growing brain is more vulnerable to long-term impact from alcohol and other drugs.







	Current Substance Use Tren Grade 12	ds
	Cigarettes Alcohol Marijuana Methamphetan	nines
60%	<u> </u>	
50%		
40%		
30%		
20%	•	
10%		
0%	<u> </u>	
2002	2004	2006

Prevalence **	2002	2004	2006	2008	20
Cigarettes	S	22% ± 3% (682)	S	25% ± 3% (800)	15% : (108
Alcohol	S	41% ± 4% (681)	S	45% ± 3% (795)	38% : (108
Marijuana	S	23% ± 3% (680)	S	30% ± 3% (795)	25% : (108
Methamphetamines	S	5% ± 2% (680)	S	4% ± 2% (393)	N/

From 2002 to 2004, change in marijuana use could not be assessed.

For More Information

Parents, schools, kids, and communities can work together to keep youth safe. To see how State and local partners are addressing these issues, see listings on the AskHYS homepage. For free guides on preventing alcohol and other drug use, call Washington State Alcohol/Drug Information Clearinghouse: 1-800-662-9111.

If you or someone you know needs help with drug abuse, contact the Washington State Alcohol/Drug Helpline at 1-800-562-1240, or online at http://www.adhl.org/

 $^{^{\}star}$ indicates a significant change (Wald $X^2,\,p{<}.05)$

^{**} Prevalence is displayed as the estimated percentage ± 95% confidence interval, with the number of respondents to this question in parentheses.

Missing Codes: S = result suppressed due to insufficient reporting from students or schools; N/P = location did not participate in the survey this year; N/G = grade not available; N/S = question was not surveyed this year; N/A = question was not asked of this grade; NB/G = this gender was not at this location